

Diagnosis and Initial Steps in the Management of PD

Miguel Situ-Kcomt, MD
Assistant Professor
Movement Disorders
Department of Neurological Sciences



1

How to Diagnose Parkinson's Disease?



- **History and Clinical Examination**
- Imaging
- Labwork?

2

History Clues

- ❖ Progressive slowness of motion observed by family or members
 - Clumsiness
 - Small handwriting
 - Shuffling gait
- ❖ Asymmetric shaking of any limb at rest
- ❖ No medications that block dopamine
- ❖ **No Red Flags**



3

Red Flags

- ❖ Rapid progression
 - Use of wheelchair or significant gait assistance in less than 5 years.
 - Significant dysphagia in less than 5 years
 - Early falls (<3 years) – More than once a year
- ❖ Severe autonomic failure
 - Urinary incontinence
 - Orthostatic Hypotension




4

Clinical Examination

➤ Bradykinesia – Slowness of motion

Plus at least 1 of these:

- Resting Tremor
- Gait problems
- Muscle rigidity (“cogwheeling”)




5

Resting Tremor



6

Gait Problems and Bradykinesia



Ianse R, Danoudis M. Freezing of Gait in Parkinson's Disease: Its Pathophysiology and Pragmatic Approaches to Management. Mov Disord Clin Pract. 2017 May-Jun; 4(3): 290-297.


7

Non-motor Symptoms

- ❖ Loss of sense of smell (anosmia)
- ❖ Difficulty Sleeping
 - REM Sleep Behavior Disorder
- ❖ Mood problems:
 - Depression
 - Anxiety
 - Apathy



8



REVIEW


CME

MDS Clinical Diagnostic Criteria for Parkinson's Disease


Ronald B. Postuma, MD, MSc,^{1†} Daniela Berg, MD,^{2†*} Matthew Stern, MD,³ Werner Poewe, MD,⁴
 C. Warren Olanow, MD, FRCPC,⁵ Wolfgang Oertel, MD,⁶ José Obeso, MD, PhD,⁷ Kenneth Marek, MD,⁸ Irene Litvan, MD,⁹
 Anthony E. Lang, OC, MD, FRCPC,¹⁰ Glenda Halliday, PhD,¹² Christopher G. Goetz, MD,¹³ Thomas Gasser, MD,²
 Bruno Dubois, MD, PhD,¹⁴ Piu Chan, MD, PhD,¹⁵ Bastiaan R. Bloem, MD, PhD,¹⁶ Charles H. Adler, MD, PhD,¹⁷
 and Günther Deuschl, MD¹⁸

¹Department of Neurology, Montreal General Hospital, Montreal, Quebec, Canada
²Department of Neurodegeneration, Hertie-Institute for Clinical Brain Research and German Center for Neurodegenerative Diseases, Tuebingen, Germany
³Penn Neurological Institute, Philadelphia, Pennsylvania, USA
⁴Department of Neurology, Innsbruck Medical University, Innsbruck, Austria
⁵Department of Neurology, The Mount Sinai Hospital, New York, New York, USA
⁶Department of Neurology, Philipps University of Marburg, Marburg, Germany
⁷University of Navarra-FIMA, Pamplona, Spain
⁸Institute for Neurodegenerative Disorders, New Haven, Connecticut, USA
⁹Department of Neurosciences, UC San Diego, La Jolla, California, USA
¹⁰Division of Neurology, Toronto Western Hospital, Toronto, Ontario, Canada
¹²Neuroscience Research Australia & University of NSW, Randwick, Australia
¹³Rush University Medical Center, Chicago, Illinois, USA
¹⁴Hopital De La Salpetriere, Paris, France
¹⁵Xuanwu Hospital of Capital of Medical University, Beijing, Peoples Republic of China
¹⁶Department of Neurology, Radboud University Medical Center, Donders Institute for Brain, Cognition and Behaviour, Nijmegen, Netherlands
¹⁷The Parkinson's Disease and Movement Disorders Center, Department of Neurology, Mayo Clinic, Scottsdale, Arizona, USA
¹⁸Department of Neurology, Christian-Albrechts University, Kiel, Germany

9



Imaging – DAT Scan



DaT SCANS WILL ONLY BE HELPFUL TO DEFINE PRIMARY FROM SECONDARY PARKINSONISMS
 THE CLINICAL CORRELATION HERE IS **MORE IMPORTANT**

10

Labwork?

There is no specific lab test that will confirm or rule out Parkinson's Disease

However, there are lab tests that may help rule out co-morbid conditions that may worsen the condition.

11

Initial Management

12

Checking labs...

- TSH
 - Hypothyroidism may contribute to slowness, fatigue, cognitive impairment
- Vitamins:
 - B1
 - B6
 - B12
 - D

13

Imaging

If there is a concern for an atypical finding:

Brain MRI with and without Contrast

- Tumors
- Strokes
- Inflammatory lesions

14

Medications

Levodopa – Converts to dopamine in the brain

Formulations:

- Carbidopa/levodopa
 - Sinemet 25/100
 - Sinemet CR 25/100 or 50/200
 - Parcopa 25/100
 - Rytary (different concentrations)



A clear and unequivocal strong response to levodopa will help with diagnosis **BUT** it does not absolutely rule out other causes

15

How to start levodopa - example

Start Sinemet 25/100 –

- 1 tablet three times a day for a week then;
- 1.5 tablets three times a day for a week then;
- 2 tablets three times a day.

Make sure to take it at around the same time every day (so for instance 8am, noon and 4pm or 8am, 2pm and 8pm)

16

Other initial medications to try



❖ Dopamine agonists:

- Pramipexole
- Ropinirole
- Rotigotine



❖ Monoamino-oxidase B inhibitors:

- Rasagiline (1mg daily)
- Selegiline (5mg twice a day)

17



UNIVERSITY OF
Nebraska
Medical Center



18